

STI management flow chart in men


STI screening schedule	
Speculum and laboratory examination	<ul style="list-style-type: none"> Monthly in all KAP (regardless of PrEP/ HIV/ ART status) If there is a risk or complaint in non-KAP
Syphilis screening	<ul style="list-style-type: none"> 3 Monthly in HIV positive KAP and PrEP client 6 monthly in HIV negative KAP annually or if there is a risk or complaint in non KAP PLHIV/ ART clients.
HIV testing	3-6 Monthly depending on the risk behavior

1st Step: only for MSM


Do risk assessment for sexual transmitted infection (only in MSM)
<ul style="list-style-type: none"> Patient's first visit to clinic Patient had an episode of discharge since the last visit Patient had unprotected sex with new partner. Partner genital signs/symptoms
If any of above questions answered "yes" → Risk assessment (+) → always treat for

2nd Step: (for all men)

Complaints of Urethral Discharge or Dysuria	
↓	
Microscopy but no discharge present (try milking)	→ <ul style="list-style-type: none"> Do urethral swab and urine test (FPU). GC (+) → treat for GC + CT Urethral smear PMNL ≥ 5 or FPU PMNL ≥ 10 → treat for CT only (add Tx for GC if no success > 1 week)



Note: Non-gonococcal urethritis is diagnosed and managed as CT



References:

- MAM STI guideline 2023